

On a theorem of Blichfeldt*

Benjamin Sambale[†]

February 17, 2024

Let G be a permutation group on $n < \infty$ objects. Let $f(g)$ be the number of fixed points of $g \in G$, and let $\{f(g) : 1 \neq g \in G\} = \{f_1, \dots, f_r\}$. In this expository note we give a character-free proof of a theorem of Blichfeldt which asserts that the order of G divides $(n - f_1) \dots (n - f_r)$. We also discuss the sharpness of this bound.

Keywords: Blichfeldt's Theorem, number of fixed points, permutation character

AMS classification: 20B05

Let us consider a permutation group G on a finite set Ω consisting of n elements. By Lagrange's Theorem applied to the symmetric group on Ω , it follows that the order $|G|$ of G is a divisor of $n!$. In order to strengthen this divisibility relation we denote the number of fixed points of a subgroup $H \leq G$ on Ω by $f(H)$. Moreover, let $f(g) := f(\langle g \rangle)$ for every $g \in G$. In 1895, Maillet [11] proved the following (see also Cameron's book [4, p. 172]).

Theorem 1 (Maillet). *Let $\{f(H) : 1 \neq H \leq G\} = \{f_1, \dots, f_r\}$. Then $|G|$ divides $(n - f_1) \dots (n - f_r)$.*

Using the newly established character theory of finite groups, Blichfeldt [1] showed in 1904 that it suffices to consider cyclic subgroups H in Maillet's Theorem (this was rediscovered by Kiyota [10]).

Theorem 2 (Blichfeldt). *Let $\{f(g) : 1 \neq g \in G\} = \{f_1, \dots, f_r\}$. Then $|G|$ divides $(n - f_1) \dots (n - f_r)$.*

For the convenience of the reader we present the elegant argument which can be found in [4, Theorem 6.5].

Proof of Blichfeldt's Theorem. Since f is the permutation character, the function ψ sending $g \in G$ to $(f(g) - f_1) \dots (f(g) - f_r)$ is a generalized character of G (i. e. a difference of ordinary complex characters). From

$$\psi(g) = \begin{cases} (n - f_1) \dots (n - f_r) & \text{if } g = 1, \\ 0 & \text{if } g \neq 1 \end{cases}$$

we conclude that ψ is a multiple of the regular character ρ of G . In particular, $\rho(1) = |G|$ divides $\psi(1) = (n - f_1) \dots (n - f_r)$. \square

It seems that no elementary proof (avoiding character theory) of Blichfeldt's Theorem has been published so far. The aim of this note is to provide such a proof.

Character-free proof of Blichfeldt's Theorem. It suffices to show that

$$\frac{1}{|G|} \sum_{g \in G} (f(g) - f_1) \dots (f(g) - f_r) \in \mathbb{Z},$$

*After the publication of the article in Expo. Math. **35** (2017), 221–225, the proof of the main theorem has been simplified.

[†]Fachbereich Mathematik, TU Kaiserslautern, 67653 Kaiserslautern, Germany, sambale@mathematik.uni-kl.de

since all summands with $g \neq 1$ vanish. Expanding the product we see that it is enough to prove

$$F_k(G) := \frac{1}{|G|} \sum_{g \in G} f(g)^k \in \mathbb{Z}$$

for $k \geq 0$. Obviously, $F_0(G) = 1$. For $k \geq 1$, note that $f(g)^k$ is the number of fixed points of G on the cartesian product Ω^k . Hence, by Burnside's lemma (often called Cauchy–Frobenius lemma, see [13]) $F_k(G)$ is the number of orbits of G on Ω^k . In particular, this is an integer. \square

It is known that Blichfeldt's Theorem can be improved by considering only the fixed point numbers of non-trivial elements of prime power order. This can be seen as follows. Let S_p be a Sylow p -subgroup of G for every prime divisor p of $|G|$. Since

$$\{f(g) : 1 \neq g \in S_p\} \subseteq \{f(g) : 1 \neq g \in G \text{ has prime power order}\} = \{f_1, \dots, f_r\},$$

Theorem 2 implies that $|S_p|$ divides $(n - f_1) \dots (n - f_r)$ for every p . Since the orders $|S_p|$ are pairwise coprime, also $|G| = \prod_p |S_p|$ is a divisor of $(n - f_1) \dots (n - f_r)$. On the other hand, it does not suffice to take the fixed point numbers of the elements of prime order. An example is given by $G = \langle (1, 2)(3, 4), (1, 3)(2, 4), (1, 2)(5, 6) \rangle$. This is a dihedral group of order 8 where every involution moves exactly four letters.

Cameron-Kiyota [5] (and independently Chillag [6]) obtained another generalization of Theorem 2 where f is assumed to be any generalized character χ of G and n is replaced by its degree $\chi(1)$. A dual version for conjugacy classes instead of characters appeared in Chillag [7].

Numerous articles addressed the question of equality in Blichfeldt's Theorem. Easy examples are given by the *regular* permutation groups. These are the transitive groups whose order coincides with the degree. In fact, by Cayley's Theorem every finite group is a regular permutation group acting on itself by multiplication. A wider class of examples consists of the *sharply k -transitive* permutation groups G for $1 \leq k \leq n$. Here, for every pair of tuples $(\alpha_1, \dots, \alpha_k), (\beta_1, \dots, \beta_k) \in \Omega^k$ with $\alpha_i \neq \alpha_j$ and $\beta_i \neq \beta_j$ for all $i \neq j$ there exists a unique $g \in G$ such that $\alpha_i^g = \beta_i$ for $i = 1, \dots, k$. Setting $\alpha_i = \beta_i$ for all i , we see that any non-trivial element of G fixes less than k points. Hence,

$$\{f(g) : 1 \neq g \in G\} \subseteq \{0, 1, \dots, k - 1\}.$$

On the other hand, if $(\alpha_1, \dots, \alpha_k)$ is fixed, then there are precisely $n(n - 1) \dots (n - k + 1)$ choices for $(\beta_1, \dots, \beta_k)$. It follows that $|G| = n(n - 1) \dots (n - k + 1)$. Therefore, we have equality in Theorem 2. Note that sharply 1-transitive and regular are the same thing. An interesting family of sharply 2-transitive groups comes from the *affine groups*

$$\text{Aff}(1, p^m) = \{\varphi : \mathbb{F}_{p^m} \rightarrow \mathbb{F}_{p^m} \mid \exists a \in \mathbb{F}_{p^m}^\times, b \in \mathbb{F}_{p^m} : \varphi(x) = ax + b \ \forall x \in \mathbb{F}_{p^m}\}$$

where \mathbb{F}_{p^m} is the field with p^m elements. More generally, all sharply 2-transitive groups are *Frobenius groups* with abelian kernel. By definition, a Frobenius group G is transitive and satisfies $\{f(g) : 1 \neq g \in G\} = \{0, 1\}$. The *kernel* K of G is the subset of fixed point free elements together with the identity. Frobenius Theorem asserts that K is a (normal) subgroup of G . For the sharply 2-transitive groups this can be proved in an elementary fashion (see [4, Exercise 1.16]), but so far no character-free proof of the full claim is known. The dihedral group $\langle (1, 2, 3, 4, 5), (2, 5)(3, 4) \rangle$ of order 10 illustrates that not every Frobenius group is sharply 2-transitive.

A typical example of a sharply 3-transitive group is $\text{SL}(2, 2^m)$ with its natural action on the set of one-dimensional subspaces of $\mathbb{F}_{2^m}^2$. We leave this claim as an exercise for the interested reader. The sharply k -transitive groups for $k \in \{2, 3\}$ were eventually classified by Zassenhaus [16, 15] using near fields (see Passman's book [14, Theorems 20.3 and 20.5]). On the other hand, there are not many sharply k -transitive groups when k is large. In fact, there is a classical theorem by Jordan [9] which was supplemented by Mathieu [12].

Theorem 3 (Jordan, Mathieu). *The sharply k -transitive permutation groups with $k \geq 4$ are given as follows:*

- (i) *the symmetric group of degree $n \geq 4$ ($k \in \{n, n - 1\}$),*
- (ii) *the alternating group of degree $n \geq 6$ ($k = n - 2$),*
- (iii) *the Mathieu group of degree 11 ($k = 4$),*

(iv) the Mathieu group of degree 12 ($k = 5$).

We remark that the Mathieu groups of degree 11 and 12 are the smallest members of the *sporadic* simple groups.

In accordance with these examples, permutation groups with equality in Theorem 2 are now called *sharp* permutation groups (this was coined by Ito-Kiyota [8]). Apart from the ones we have already seen, there are more examples. For instance, the symmetry group of a square acting on the four vertices has order 8 (again a dihedral group) and the non-trivial fixed point numbers are 0 and 2. Recently, Brozovic [3] gave a description of the primitive sharp permutation groups G such that $\{f(g) : 1 \neq g \in G\} = \{0, k\}$ for some $k \geq 1$. Here, a permutation group is *primitive* if it is transitive and any one-point stabilizer is a maximal subgroup. The complete classification of the sharp permutation groups is widely open.

Finally, we use the opportunity to mention a related result by Bochert [2] where the divisibility relation of $|G|$ is replaced by an inequality. As usual $\lfloor x \rfloor$ denotes the largest integer less than or equal to $x \in \mathbb{R}$.

Theorem 4 (Bochert). *If G is primitive, then $|G| \leq n(n-1)\dots(n - \lfloor n/2 \rfloor + 1)$ unless G is the symmetric group or the alternating group of degree n .*

Acknowledgment

This work is supported by the German Research Foundation (project SA 2864/1-1) and the Daimler and Benz Foundation (project 32-08/13).

References

- [1] H. F. Blichfeldt, *A theorem concerning the invariants of linear homogeneous groups, with some applications to substitution-groups*, Trans. Amer. Math. Soc. **5** (1904), 461–466.
- [2] A. Bochert, *Ueber die Zahl der verschiedenen Werthe, die eine Function gegebener Buchstaben durch Vertauschung derselben erlangen kann*, Math. Ann. **33** (1889), 584–590.
- [3] D. P. Brozovic, *The classification of primitive sharp permutation groups of type $\{0, k\}$* , Comm. Algebra **42** (2014), 3028–3062.
- [4] P. J. Cameron, *Permutation groups*, London Mathematical Society Student Texts, Vol. 45, Cambridge University Press, Cambridge, 1999.
- [5] P. J. Cameron and M. Kiyota, *Sharp characters of finite groups*, J. Algebra **115** (1988), 125–143.
- [6] D. Chillag, *Character values of finite groups as eigenvalues of nonnegative integer matrices*, Proc. Amer. Math. Soc. **97** (1986), 565–567.
- [7] D. Chillag, *On a congruence of Blichfeldt concerning the order of finite groups*, Proc. Amer. Math. Soc. **136** (2008), 1961–1966.
- [8] T. Ito and M. Kiyota, *Sharp permutation groups*, J. Math. Soc. Japan **33** (1981), 435–444.
- [9] C. Jordan, *Sur la limite de transitivité des groupes non-alternées*, Bull. Soc. Math. France **1** (1873), 40–71.
- [10] M. Kiyota, *An inequality for finite permutation groups*, J. Combin. Theory Ser. A **27** (1979), 119.
- [11] E. Maillet, *Sur quelques propriétés des groupes de substitutions d'ordre donné*, Ann. Fac. Sci. Toulouse Sci. Math. Sci. Phys. **9** (1895), 1–22.
- [12] É. Mathieu, *Mémoire sur l'étude des fonctions de plusieurs quantités, sur la manière de les former et sur les substitutions qui les laissent invariables*, J. Math. Pures Appl. **6** (1861), 241–323.
- [13] P. M. Neumann, *A lemma that is not Burnside's*, Math. Sci. **4** (1979), 133–141.

- [14] D. S. Passman, *Permutation groups*, Dover Publications, Inc., Mineola, N.Y., 2012 (revised reprint of the 1968 original).
- [15] H. Zassenhaus, *Kennzeichnung endlicher linearer Gruppen als Permutationsgruppen*, Abh. Math. Sem. Univ. Hamburg **11** (1935), 17–40.
- [16] H. Zassenhaus, *Über endliche Fastkörper*, Abh. Math. Sem. Univ. Hamburg **11** (1935), 187–220.