

Real characters in nilpotent blocks

Benjamin Sambale*

February 5, 2023

Dedicated to Pham Huu Tiep on the occasion of his 60th birthday.

Abstract

We prove that the number of irreducible real characters in a nilpotent block of a finite group is locally determined. We further conjecture that the Frobenius–Schur indicators of those characters can be computed for $p = 2$ in terms of the extended defect group. We derive this from a more general conjecture on the Frobenius–Schur indicator of projective indecomposable characters of 2-blocks with one simple module. This extends results of Murray on 2-blocks with cyclic and dihedral defect groups.

Keywords: real characters; Frobenius–Schur indicators; nilpotent blocks

AMS classification: 20C15, 20C20

1 Introduction

An important task in representation theory is to determine global invariants of a finite group G by means of local subgroups. Dade’s conjecture, for instance, predicts the number of irreducible characters $\chi \in \text{Irr}(G)$ such that the p -part $\chi(1)_p$ is a given power of a prime p (see [23, Conjecture 9.25]). Since Gow’s work [7], there has been an increasing interest in counting real (i. e. real-valued) characters and more generally characters with a given field of values.

The quaternion group Q_8 testifies that a real irreducible character χ is not always afforded by a representation over the real numbers. The precise behavior is encoded by the *Frobenius–Schur indicator* (F-S indicator, for short)

$$\epsilon(\chi) := \frac{1}{|G|} \sum_{g \in G} \chi(g^2) = \begin{cases} 0 & \text{if } \bar{\chi} \neq \chi, \\ 1 & \text{if } \chi \text{ is realized by a real representation,} \\ -1 & \text{if } \chi \text{ is real, but not realized by a real representation.} \end{cases} \quad (1)$$

A new interpretation of the F-S indicator in terms of superalgebras has been given recently in [13]. The case of the dihedral group D_8 shows that $\epsilon(\chi)$ is not determined by the character table of G . The computation of F-S indicators can be a surprisingly difficult task, which has not been fully completed

*Institut für Algebra, Zahlentheorie und Diskrete Mathematik, Leibniz Universität Hannover, Welfengarten 1, 30167 Hannover, Germany, sambale@math.uni-hannover.de

for the simple groups of Lie type, for instance (see [25]). Problem 14 on Brauer's famous list [2] asks for a group-theoretical interpretation of the number of $\chi \in \text{Irr}(G)$ with $\epsilon(\chi) = 1$.

To obtain deeper insights, we fix a prime p and assume that χ lies in a p -block B of G with defect group D . By complex conjugation we obtain another block \overline{B} of G . If $\overline{B} \neq B$, then clearly $\epsilon(\chi) = 0$ for all $\chi \in \text{Irr}(B)$. Hence, we assume that B is real, i.e. $\overline{B} = B$. John Murray [18, 19] has computed the F-S indicators when D is a cyclic 2-group or a dihedral 2-group (including the Klein four-group). His results depend on the fusion system of B , on Erdmann's classification of tame blocks and on the structure of the so-called *extended defect group* E of B (see Definition 7 below). For $p > 2$ and D cyclic, he obtained in [20] partial information on the F-S indicators in terms of the Brauer tree of B .

The starting point of my investigation is the well-known fact that 2-blocks with cyclic defect groups are nilpotent. Assume that B is nilpotent and real. If B is the principal block, then $G = \text{O}_{p'}(G)D$ and $\text{Irr}(B) = \text{Irr}(G/\text{O}_{p'}(G)) = \text{Irr}(D)$. In this case the F-S indicators of B are determined by D alone. Thus, suppose that B is non-principal. By Broué–Puig [4], there exists a height-preserving bijection $\text{Irr}(D) \rightarrow \text{Irr}(B)$, $\lambda \mapsto \lambda * \chi_0$ where $\chi_0 \in \text{Irr}(B)$ is a fixed character of height 0 (see also [16, Definition 8.10.2]). However, this bijection does not in general preserve F-S indicators. For instance, the dihedral group D_{24} has a nilpotent 2-block with defect group C_4 and a nilpotent 3-block with defect group C_3 , although every character of D_{24} is real. Our main theorem asserts that the number of real characters in a nilpotent block is nevertheless locally determined. To state it, we introduce the *extended inertial group*

$$\text{N}_G(D, b_D)^* := \{g \in \text{N}_G(D) : b_D^g \in \{b_D, \overline{b_D}\}\}$$

where b_D is a Brauer correspondent of B in $\text{DC}_G(D)$.

Theorem A. *Let B be a real, nilpotent p -block of a finite group G with defect group D . Let b_D be a Brauer correspondent of B in $\text{DC}_G(D)$. Then the number of real characters in $\text{Irr}(B)$ of height h coincides with the number of characters $\lambda \in \text{Irr}(D)$ of degree p^h such that $\lambda^t = \overline{\lambda}$ where*

$$\text{N}_G(D, b_D)^*/\text{DC}_G(D) = \langle t\text{DC}_G(D) \rangle.$$

If $p > 2$, then all real characters in $\text{Irr}(B)$ have the same F-S indicator.

In contrast to arbitrary blocks, Theorem A implies that nilpotent real blocks have at least one real character (cf. [20, p. 92] and [8, Theorem 5.3]). If $\overline{b_D} = b_D$, then B and D have the same number of real characters, because $\text{N}_G(D, b_D) = \text{DC}_G(D)$. This recovers a result of Murray [18, Lemma 2.2]. As another consequence, we will derive in Proposition 5 a real version of Eaton's conjecture [5] for nilpotent blocks as put forward by Héthelyi–Horváth–Szabó [12].

The F-S indicators of real characters in nilpotent blocks seem to lie somewhat deeper. We still conjecture that they are locally determined by a defect pair (see Definition 7) for $p = 2$ as follows.

Conjecture B. *Let B be a real, nilpotent, non-principal 2-block of a finite group G with defect pair (D, E) . Then there exists a height preserving bijection $\Gamma : \text{Irr}(D) \rightarrow \text{Irr}(B)$ such that*

$$\epsilon(\Gamma(\lambda)) = \frac{1}{|D|} \sum_{e \in E \setminus D} \lambda(e^2) \tag{2}$$

for all $\lambda \in \text{Irr}(D)$.

The right hand side of (2) was introduced and studied by Gow [8, Lemma 2.1] more generally for any groups $D \leq E$ with $|E : D| = 2$. This invariant was later coined the *Gow indicator* by Murray [20, Eq. (2)]. For 2-blocks of defect 0, Conjecture B confirms the known fact that real characters of 2-defect 0 have F-S indicator 1 (see [8, Theorem 5.1]). There is no such result for odd primes p . As a matter of fact, every real character has p -defect 0 whenever p does not divide $|G|$. In Theorem 10 we prove Conjecture B for abelian defect groups D . Then it also holds for all quasisimple groups G by work of An–Eaton [1]. Murray’s results mentioned above, imply Conjecture B also for dihedral D .

For $p > 2$, the common F-S indicator in the situation of Theorem A is not locally determined. For instance, $G = Q_8 \rtimes C_9 = \text{SmallGroup}(72, 3)$ has a non-principal real 3-block with $D \cong C_9$ and common F-S indicator -1 , while its Brauer correspondent in $N_G(D) \cong C_{18}$ has common F-S indicator 1. Nevertheless, for cyclic defect groups D we find another way to compute this F-S indicator in Theorem 3 below.

Our second conjecture applies more generally to blocks with only one simple module.

Conjecture C. *Let B be a real, non-principal 2-block with defect pair (D, E) and a unique projective indecomposable character Φ . Then*

$$\epsilon(\Phi) = |\{x \in E \setminus D : x^2 = 1\}|.$$

Here $\epsilon(\Phi)$ is defined by extending (1) linearly. If $\epsilon(\Phi) = 0$, then E does not split over D and Conjecture C holds (see Proposition 8 below). Conjecture C implies a stronger, but more technical statement on 2-blocks with a Brauer correspondent with one simple module (see Theorem 13 below). This allows us to prove the following.

Theorem D. *Conjecture C implies Conjecture B.*

We remark that our proof of Theorem D does not work block-by-block. For solvable groups we offer a purely group-theoretical version of Conjecture C at the end of Section 4.

Theorem E. *Conjectures B and C hold for all nilpotent 2-blocks of solvable groups.*

We have checked Conjectures B and C with GAP [6] in many examples using the libraries of small groups, perfect groups and primitive groups.

2 Theorem A and its consequences

Our notation follows closely Navarro’s book [22]. In particular, G^0 denotes the set of p -regular elements of a finite group G . Let B be a p -block of G with defect group D . Recall that a B -*subsection* is a pair (u, b) where $u \in D$ and b is a Brauer correspondent of B in $C_G(u)$. For $\chi \in \text{Irr}(B)$ and $\varphi \in \text{IBr}(b)$ we denote the corresponding generalized decomposition number by $d_{\chi\varphi}^u$. If $u = 1$, we obtain the (ordinary) decomposition number $d_{\chi\varphi} = d_{\chi\varphi}^1$. We put $l(b) = |\text{IBr}(b)|$ as usual.

Following [22, p. 114], we define a class function $\chi^{(u,b)}$ by

$$\chi^{(u,b)}(us) := \sum_{\varphi \in \text{IBr}(b)} d_{\chi\varphi}^u \varphi(s)$$

for $s \in C_G(u)^0$ and $\chi^{(u,b)}(x) = 0$ whenever x is outside the p -section of u . If \mathcal{R} is a set of representatives for the G -conjugacy classes of B -subsections, then $\chi = \sum_{(u,b) \in \mathcal{R}} \chi^{(u,b)}$ by Brauer's second main theorem (see [22, Problem 5.3]). Now suppose that B is nilpotent and $\lambda \in \text{Irr}(D)$. By [16, Proposition 8.11.4], each Brauer correspondent b of B is nilpotent and in particular $l(b) = 1$. Broué–Puig [4] have shown that, if χ has height 0, then

$$\lambda * \chi := \sum_{(u,b) \in \mathcal{R}} \lambda(u) \chi^{(u,b)} \in \text{Irr}(B)$$

and $(\lambda * \chi)(1) = \lambda(1)\chi(1)$. Note also that $d_{\lambda * \chi, \varphi}^u = \lambda(u)d_{\chi, \varphi}^u$.

Proof of Theorem A. Let \mathcal{R} be a set of representatives for the G -conjugacy classes of B -subsections $(u, b_u) \leq (D, b_B)$ (see [22, p. 219]). Since B is nilpotent, we have $\text{IBr}(b_u) = \{\varphi_u\}$ for all $(u, b_u) \in \mathcal{R}$. Since the Brauer correspondence is compatible with complex conjugation, $(u, \overline{b_u})^t \leq (D, \overline{b_D})^t = (D, b_D)$ where $\text{N}_G(D, b_D)^* / \text{DC}_G(D) = \langle t\text{DC}_G(D) \rangle$. Thus, $(u, \overline{b_u})^t$ is D -conjugate to some $(u', b_{u'}) \in \mathcal{R}$.

If $p > 2$, there exists a unique p -rational character $\chi_0 \in \text{Irr}(B)$ of height 0, which must be real by uniqueness (see [4, Remark after Theorem 1.2]). If $p = 2$, there is a 2-rational real character $\chi_0 \in \text{Irr}(B)$ of height 0 by [8, Theorem 5.1]. Then $d_{\chi_0, \varphi_u}^u = d_{\chi_0, \overline{\varphi_u}}^u \in \mathbb{Z}$ and

$$\overline{\chi_0^{(u, b_u)}} = \chi_0^{(u, \overline{b_u})} = \chi_0^{(u, \overline{b_u})^t} = \chi_0^{(u', b_{u'})}.$$

Now let $\lambda \in \text{Irr}(D)$. Then

$$\overline{\lambda * \chi_0} = \sum_{(u, b_u) \in \mathcal{R}} \overline{\lambda(u) \chi_0^{(u, b_u)}} = \sum_{(u, b_u) \in \mathcal{R}} \overline{\lambda(u)} \overline{\chi_0^{(u, b_u)}} = \sum_{(u, b_u) \in \mathcal{R}} \overline{\lambda(u)} \chi_0^{(u', b_{u'})}.$$

Since the class functions $\chi_0^{(u, b)}$ have disjoint support, they are linearly independent. Therefore, $\lambda * \chi_0$ is real if and only if $\lambda(u^t) = \lambda(u') = \overline{\lambda(u)}$ for all $(u, b_u) \in \mathcal{R}$. Since every conjugacy class of D is represented by some u with $(u, b_u) \in \mathcal{R}$, we conclude that $\lambda * \chi_0$ is real if and only if $\lambda^t = \overline{\lambda}$. Moreover, if $\lambda(1) = p^h$, then $\lambda * \chi_0$ has height h . This proves the first claim.

To prove the second claim, let $p > 2$ and $\text{IBr}(B) = \{\varphi\}$. Then the decomposition numbers $d_{\lambda * \chi_0, \varphi} = \lambda(1)$ are powers of p ; in particular they are odd. A theorem of Thompson and Willems (see [26, Theorem 2.8]) states that all real characters χ with $d_{\chi, \varphi}$ odd have the same F-S indicator. So in our situation all real characters in $\text{Irr}(B)$ have the same F-S indicator. \square

Since the automorphism group of a p -group is ‘‘almost always’’ a p -group (see [11]), the following consequence is of interest.

Corollary 1. *Let B be a real, nilpotent p -block with defect group D such that p and $|\text{Aut}(D)|$ are odd. Then B has a unique real character.*

Proof. The hypothesis on $\text{Aut}(D)$ implies that $\text{N}_G(D, b_D)^* = \text{DC}_G(D)$. Hence by Theorem A, the number of real characters in $\text{Irr}(B)$ is the number of real characters in D . Since $p > 2$, the trivial character is the only real character of D . \square

The next lemma is a consequence of Brauer's second main theorem and the fact that $|\{g \in G : g^2 = x\}| = |\{g \in C_G(x) : g^2 = x\}|$ is locally determined for $g, x \in G$.

Lemma 2 (Brauer). *For every p -block B of G and every B -subsection (u, b) with $\varphi \in \text{IBr}(b)$ we have*

$$\sum_{\chi \in \text{Irr}(B)} \epsilon(\chi) d_{\chi\varphi}^u = \sum_{\psi \in \text{Irr}(b)} \epsilon(\psi) d_{\psi\varphi}^u = \sum_{\psi \in \text{Irr}(b)} \epsilon(\psi) \frac{\psi(u)}{\psi(1)} d_{\psi\varphi}.$$

If $l(b) = 1$, then

$$\sum_{\chi \in \text{Irr}(B)} \epsilon(\chi) d_{\chi\varphi}^u = \frac{1}{\varphi(1)} \sum_{\psi \in \text{Irr}(b)} \epsilon(\psi) \psi(u).$$

Proof. The first equality is [3, Theorem 4A]. The second follows from $u \in Z(C_G(u))$. If $l(b) = 1$, then $\psi(1) = d_{\psi\varphi}\varphi(1)$ for $\psi \in \text{Irr}(b)$ and the last claim follows. \square

Recall that a *canonical* character of B is a character $\theta \in \text{Irr}(DC_G(D))$ lying in a Brauer correspondent of B such that $D \leq \text{Ker}(\theta)$ (see [22, Theorem 9.12]). We define the *extended stabilizer*

$$N_G(D)_\theta^* := \{g \in N_G(D) : \theta^g \in \{\theta, \bar{\theta}\}\}.$$

The following results adds some detail to the nilpotent case of [20, Theorem 1].

Theorem 3. *Let B be a real, nilpotent p -block with cyclic defect group $D = \langle u \rangle$ and $p > 2$. Let $\theta \in \text{Irr}(C_G(D))$ be a canonical character of B and set $T := N_G(D)_\theta^*$. Then one of the following holds:*

- (1) $\bar{\theta} \neq \theta$. *All characters in $\text{Irr}(B)$ are real with F-S indicator $\epsilon(\theta^T)$.*
- (2) $\bar{\theta} = \theta$. *The unique non-exceptional character $\chi_0 \in \text{Irr}(B)$ is the only real character in $\text{Irr}(B)$ and $\epsilon(\chi_0) = \text{sgn}(\chi_0(u))\epsilon(\theta)$ where $\text{sgn}(\chi_0(u))$ is the sign of $\chi_0(u)$.*

Proof. Let b_D be a Brauer correspondent of B in $C_G(D)$ containing θ . Then $T = N_G(D, b_D)^*$. If $\bar{\theta} \neq \theta$, then T inverts the elements of D since $p > 2$. Thus, Theorem A implies that all characters in $\text{Irr}(B)$ are real. By [20, Theorem 1(v)], the common F-S indicator is the Gow indicator of θ with respect to T . This is easily seen to be $\epsilon(\theta^T)$ (see [20, after Eq. (2)]).

Now assume that $\bar{\theta} = \theta$. Here Theorem A implies that the unique p -rational character $\chi_0 \in \text{Irr}(B)$ is the only real character. In particular, χ_0 must be the unique non-exceptional character. Note that (u, b_D) is a B -subsection and $\text{IBr}(b_D) = \{\varphi\}$. Since χ_0 is p -rational, $d_{\chi_0\varphi}^u = \pm 1$. Since all Brauer correspondents of B in $C_G(u)$ are conjugate under $N_G(D)$, the generalized decomposition numbers are Galois conjugate, in particular $d_{\chi_0\varphi}^u$ does not depend on the choice of b_D . Hence,

$$\chi_0(u) = |N_G(D) : N_G(D)_\theta| d_{\chi_0\varphi}^u \varphi(1)$$

and $d_{\chi_0\varphi}^u = \text{sgn}(\chi_0(u))$. Moreover, θ is the unique non-exceptional character of b_D and $\theta(u) = \theta(1)$. By Lemma 2, we obtain

$$\epsilon(\chi_0) = \text{sgn}(\chi_0(u)) \sum_{\chi \in \text{Irr}(B)} \epsilon(\chi) d_{\chi\varphi}^u = \frac{\text{sgn}(\chi_0(u))}{\varphi(1)} \sum_{\psi \in \text{Irr}(b_D)} \epsilon(\psi) \psi(u) = \text{sgn}(\chi_0(u))\epsilon(\theta). \quad \square$$

If B is a nilpotent block with canonical character $\theta \neq \bar{\theta}$, the common F-S indicator of the real characters in $\text{Irr}(B)$ is not always $\epsilon(\theta^T)$ as in Theorem 3. A counterexample is given by a certain 3-block of $G = \text{SmallGroup}(288, 924)$ with defect group $D \cong C_3 \times C_3$.

We now restrict ourselves to 2-blocks. Héthelyi–Horváth–Szabó [12] introduced four conjectures, which are real versions of Brauer’s conjecture, Olsson’s conjecture and Eaton’s conjecture. We only state the strongest of them, which implies the remaining three. Let $D^{(0)} := D$ and $D^{(k+1)} := [D^{(k)}, D^{(k)}]$ for $k \geq 0$ be the members of the derived series of D .

Conjecture 4 (Héthelyi–Horváth–Szabó). *Let B be a 2-block with defect group D . For every $h \geq 0$, the number of real characters in $\text{Irr}(B)$ of height $\leq h$ is bounded by the number of elements of $D/D^{(h+1)}$ which are real in $N_G(D)/D^{(h+1)}$.*

A conjugacy class K of G is called *real* if $K = K^{-1} := \{x^{-1} : x \in K\}$. A conjugacy class K of a normal subgroup $N \trianglelefteq G$ is called *real under G* if there exists $g \in G$ such that $K^g = K^{-1}$.

Proposition 5. *Let B be a nilpotent 2-block with defect group D and Brauer correspondent b_D in $DC_G(D)$. Then the number of real characters in $\text{Irr}(B)$ of height $\leq h$ is bounded by the number of conjugacy classes of $D/D^{(h+1)}$ which are real under $N_G(D, b_D)^*/D^{(h+1)}$. In particular, Conjecture 4 holds for B .*

Proof. We may assume that B is real. As in the proof of Theorem A, we fix some 2-rational real character $\chi_0 \in \text{Irr}(B)$ of height 0. Now $\lambda * \chi_0$ has height $\leq h$ if and only if $\lambda(1) \leq p^h$ for $\lambda \in \text{Irr}(B)$. By [14, Theorem 5.12], the characters of degree $\leq p^h$ in $\text{Irr}(D)$ lie in $\text{Irr}(D/D^{(h+1)})$. By Theorem A, $\lambda * \chi_0$ is real if and only if $\lambda^t = \bar{\lambda}$. By Brauer’s permutation lemma (see [23, Theorem 2.3]), the number of those characters λ coincides with the number of conjugacy classes K of $D/D^{(h+1)}$ such that $K^t = K^{-1}$. Now Conjecture 4 follows from $N_G(D, b_D)^* \leq N_G(D)$. \square

3 Extended defect groups

We continue to assume that $p = 2$. As usual we choose a complete discrete valuation ring \mathcal{O} such that $F := \mathcal{O}/J(\mathcal{O})$ is an algebraically closed field of characteristic 2. Let $\text{Cl}(G)$ be the set of conjugacy classes of G . For $K \in \text{Cl}(G)$ let $K^+ := \sum_{x \in K} x \in Z(FG)$ be the class sum of K . We fix a 2-block B of FG with block idempotent $1_B = \sum_{K \in \text{Cl}(G)} a_K K^+$ where $a_K \in F$. The central character of B is defined by

$$\lambda_B : Z(FG) \rightarrow F, \quad K^+ \mapsto \left(\frac{|K|\chi(g)}{\chi(1)} \right)^*$$

where $g \in K$, $\chi \in \text{Irr}(B)$ and $*$ denotes the canonical reduction $\mathcal{O} \rightarrow F$ (see [22, Chapter 2]).

Since $\lambda_B(1_B) = 1$, there exists $K \in \text{Cl}(G)$ such that $a_K \neq 0 \neq \lambda_B(K^+)$. We call K a *defect class* of B . By [22, Corollary 3.8], K consists of elements of odd order. According to [22, Corollary 4.5], a Sylow 2-subgroup D of $C_G(x)$ where $x \in K$ is a defect group of B . For $x \in K$ let

$$C_G(x)^* := \{g \in G : gxg^{-1} = x^{\pm 1}\} \leq G$$

be the *extended centralizer* of x .

Proposition 6 (Gow, Murray). *Every real 2-block B has a real defect class K . Let $x \in K$. Choose a Sylow 2-subgroup E of $C_G(x)^*$ and put $D := E \cap C_G(x)$. Then the G -conjugacy class of the pair (D, E) does not depend on the choice of K or x .*

Proof. For the principal block (which is always real since it contains the trivial character), $K = \{1\}$ is a real defect class and $E = D$ is a Sylow 2-subgroup of G . Hence, the uniqueness follows from Sylow’s theorem. Now suppose that B is non-principal. The existence of K was first shown in [8, Theorem 5.5]. Let L be another real defect class of B and choose $y \in L$. By [9, Corollary 2.2], we may assume after conjugation that E is also a Sylow 2-subgroup of $C_G(y)^*$. Let $D_x := E \cap C_G(x)$ and $D_y := E \cap C_G(y)$. We may assume that $|E : D_x| = 2 = |E : D_y|$ (cf. the remark after the proof).

We now introduce some notation in order to apply [17, Proposition 14]. Let $\Sigma = \langle \sigma \rangle \cong C_2$. We consider FG as an $F[G \times \Sigma]$ -module where G acts by conjugation and $g^\sigma = g^{-1}$ for $g \in G$ (observe that these actions indeed commute). For $H \leq G \times \Sigma$ let

$$\mathrm{Tr}_H^{G \times \Sigma} : (FG)^H \rightarrow (FG)^{G \times \Sigma}, \alpha \mapsto \sum_{x \in \mathcal{R}} \alpha^x$$

be the *relative trace* with respect to H , where \mathcal{R} denotes a set of representatives of the right cosets of H in $G \times \Sigma$. By [17, Proposition 14], we have $1_B \in \mathrm{Tr}_{E_x}^{G \times \Sigma}(FG)$ where $E_x := D_x \langle e_x \sigma \rangle$ for some $e_x \in E \setminus D_x$. By the same result we also obtain that $D_y \langle e_y \sigma \rangle$ with $e_y \in E \setminus D_y$ is G -conjugate to E_x . This implies that D_y is conjugate to D_x inside $N_G(E)$. In particular, (D_x, E) and (D_y, E) are G -conjugate as desired. \square

Definition 7. In the situation of Proposition 6 we call E an *extended defect group* and (D, E) a *defect pair* of B .

We stress that real 2-blocks can have non-real defect classes and non-real blocks can have real defect classes (see [10, Theorem 3.5]).

It is easy to show that non-principal real 2-blocks cannot have maximal defect (see [22, Problem 3.8]). In particular, the trivial class cannot be a defect class and consequently, $|E : D| = 2$ in those cases. For non-real blocks we define the extended defect group by $E := D$ for convenience. Every given pair of 2-groups $D \leq E$ with $|E : D| = 2$ occurs as a defect pair of a real (nilpotent) block. To see this, let $Q \cong C_3$ and $G = Q \rtimes E$ with $C_E(Q) = D$. Then G has a unique non-principal block with defect pair (D, E) .

We recall from [14, p. 49] that

$$\sum_{\chi \in \mathrm{Irr}(G)} \epsilon(\chi) \chi(g) = |\{x \in G : x^2 = g\}| \quad (3)$$

for all $g \in G$. The following proposition provides some interesting properties of defect pairs.

Proposition 8 (Gow, Murray). *Let B be a real 2-block with defect pair (D, E) . Let b_D be a Brauer correspondent of B in $DC_G(D)$. Then the following holds:*

- (i) $N_G(D, b_D)^* = N_G(D, b_D)E$. In particular, b_D is real if and only if $E = DC_E(D)$.
- (ii) For $u \in D$, we have $\sum_{\chi \in \mathrm{Irr}(B)} \epsilon(\chi) \chi(u) \geq 0$ with strict inequality if and only if u is G -conjugate to e^2 for some $e \in E \setminus D$. In particular, E splits over D if and only if $\sum_{\chi \in \mathrm{Irr}(B)} \epsilon(\chi) \chi(1) > 0$.
- (iii) E/D' splits over D/D' if and only if all height zero characters in $\mathrm{Irr}(B)$ have non-negative F - S indicator.

Proof.

- (i) See [19, Lemma 1.8] and [18, Theorem 1.4].
- (ii) See [19, Lemma 1.3].
- (iii) See [8, Theorem 5.6]. \square

The next proposition extends [18, Lemma 1.3].

Corollary 9. *Suppose that B is a 2-block with defect pair (D, E) where D is abelian. Then E splits over D if and only if all characters in $\text{Irr}(B)$ have non-negative F-S indicator.*

Proof. If B is non-real, then $E = D$ splits over D and all characters in $\text{Irr}(B)$ have F-S indicator 0. Hence, let $\overline{B} = B$. By Kessar–Malle [15], all characters in $\text{Irr}(B)$ have height 0. Hence, the claim follows from Proposition 8(iii). \square

Theorem 10. *Let B be a real, nilpotent 2-block with defect pair (D, E) where D is abelian. If E splits over D , then all real characters in $\text{Irr}(B)$ have F-S indicator 1. Otherwise exactly half of the real characters have F-S indicator 1. In either case, Conjecture B holds for B .*

Proof. If E splits over D , then all real characters in $\text{Irr}(B)$ have F-S indicator 1 by Corollary 9. Otherwise we have $\sum_{\chi \in \text{Irr}(B)} \epsilon(\chi) = 0$ by Proposition 8(ii), because all characters in $\text{Irr}(B)$ have the same degree. Hence, exactly half of the real characters have F-S indicator 1. Using Theorem A we can determine the number of characters for each F-S indicator. For the last claim, we may therefore replace B by the unique non-principal block of $G = Q \rtimes E$ where $Q \cong C_3$ and $C_E(Q) = D$ (mentioned above). In this case Conjecture B follows from Gow [8, Lemma 2.2] or Theorem E. \square

Example 11. Let B be a real block with defect group $D \cong C_4 \times C_2$. Then B is nilpotent since $\text{Aut}(D)$ is a 2-group and D is abelian. Moreover $|\text{Irr}(B)| = 8$. The F-S indicators depend not only on E , but also on the way D embeds into E . The following cases can occur (here M_{16} denotes the modular group and [16, 3] refers to the small group library):

F-S indicators	E
+++++	$D_8 \times C_2$
++++--	$Q_8 \times C_2, C_4 \rtimes C_4$ with $\Phi(D) = E'$
++++ 0 0 0 0	$D, D \times C_2, D_8 * C_4, [16, 3]$
++-- 0 0 0 0	$C_4^2, C_8 \times C_2, M_{16}, C_4 \rtimes C_4$ with $\Phi(D) \neq E'$

The F-S indicator $\epsilon(\Phi)$ appearing in Conjecture C has an interesting interpretation as follows. Let $\Omega := \{g \in G : g^2 = 1\}$. The conjugation action of G on Ω turns $F\Omega$ into an FG -module, called the *involution module*.

Lemma 12 (Murray). *Let B be a real 2-block and $\varphi \in \text{IBr}(B)$. Then $\epsilon(\Phi_\varphi)$ is the multiplicity of φ as a constituent of the Brauer character of $F\Omega$.*

Proof. See [18, Lemma 2.6]. \square

Next we develop a local version of Conjecture C. Let B be a real 2-block with defect pair (D, E) and B -subsection (u, b) . If $E = DC_E(u)$, then b is real and $(C_D(u), C_E(u))$ is a defect pair of b by [19, Lemma 2.6] applied to the subpair $(\langle u \rangle, b)$. Conversely, if b is real, we may assume that $(C_D(u), C_E(u))$ is a defect pair of b by [19, Theorem 2.7]. If b is non-real, we may assume that $(C_D(u), C_D(u)) = (C_D(u), C_E(u))$ is a defect pair of b .

Theorem 13. *Let B be 2-block of a finite group G with defect pair (D, E) . Suppose that Conjecture C holds for all Brauer correspondents of B in sections of G . Let (u, b) be a B -subsection with defect pair $(C_D(u), C_E(u))$ such that $\text{IBr}(b) = \{\varphi\}$. Then*

$$\sum_{\chi \in \text{Irr}(B)} \epsilon(\chi) d_{\chi\varphi}^u = \begin{cases} |\{x \in D : x^2 = u\}| & \text{if } B \text{ is the principal block,} \\ |\{x \in E \setminus D : x^2 = u\}| & \text{otherwise.} \end{cases}$$

Proof. If B is not real, then B is non-principal and $E = D$. It follows that $\epsilon(\chi) = 0$ for all $\chi \in \text{Irr}(B)$ and

$$|\{x \in E \setminus D : x^2 = u\}| = 0.$$

Hence, we may assume that B is real. By Lemma 2, we have

$$\sum_{\chi \in \text{Irr}(B)} \epsilon(\chi) d_{\chi\varphi}^u = \sum_{\psi \in \text{Irr}(b)} \epsilon(\psi) d_{\psi\varphi}^u = \frac{1}{\varphi(1)} \sum_{\psi \in \text{Irr}(b)} \epsilon(\psi) \psi(u). \quad (4)$$

Suppose that B is the principal block. Then b is the principal block of $C_G(u)$ by Brauer's third main theorem (see [22, Theorem 6.7]). The hypothesis $l(b) = 1$ implies that $\varphi = 1_{C_G(u)}$ and $C_G(u)$ has a normal 2-complement N (see [22, Corollary 6.13]). It follows that $\text{Irr}(b) = \text{Irr}(C_G(u)/N) = \text{Irr}(C_D(u))$ and

$$\sum_{\psi \in \text{Irr}(b)} \epsilon(\psi) d_{\psi\varphi}^u = \sum_{\lambda \in \text{Irr}(C_D(u))} \epsilon(\lambda) \lambda(u) = |\{x \in C_D(u) : x^2 = u\}|$$

by (3). Since every $x \in D$ with $x^2 = u$ lies in $C_D(u)$, we are done in this case.

Now let B be a non-principal real 2-block. If b is not real, then (4) shows that $\sum_{\chi \in \text{Irr}(B)} \epsilon(\chi) d_{\chi\varphi}^u = 0$. On the other hand, we have $C_E(u) = C_D(u) \leq D$ and $|\{x \in E \setminus D : x^2 = u\}| = 0$. Hence, we may assume that b is real. Since every $x \in E$ with $x^2 = u$ lies in $C_E(u)$, we may assume that $u \in Z(G)$ by (4).

Then $\chi(u) = d_{\chi\varphi}^u \varphi(1)$ for all $\chi \in \text{Irr}(B)$. If $u^2 \notin \text{Ker}(\chi)$, then $\chi(u) \notin \mathbb{R}$ and $\epsilon(\chi) = 0$. Thus, it suffices to sum over χ with $d_{\chi\varphi}^u = \pm d_{\chi\varphi}$. Let $Z := \langle u \rangle \leq Z(G)$ and $\bar{G} := G/Z$. Let \hat{B} be the unique (real) block of \bar{G} dominated by B . By [19, Lemma 1.7], (\bar{D}, \bar{E}) is a defect pair for \hat{B} . Then, using [14, Lemma 4.7] and Conjecture C for B and \hat{B} , we obtain

$$\begin{aligned} \sum_{\chi \in \text{Irr}(B)} \epsilon(\chi) d_{\chi\varphi}^u &= \sum_{\chi \in \text{Irr}(B)} \epsilon(\chi) (d_{\chi\varphi} + d_{\chi\varphi}^u) - \sum_{\chi \in \text{Irr}(B)} \epsilon(\chi) d_{\chi\varphi} \\ &= 2 \sum_{\chi \in \text{Irr}(\hat{B})} \epsilon(\chi) d_{\chi\varphi} - \sum_{\chi \in \text{Irr}(B)} \epsilon(\chi) d_{\chi\varphi} \\ &= 2|\{\bar{x} \in \bar{E} \setminus \bar{D} : \bar{x}^2 = 1\}| - |\{x \in E \setminus D : x^2 = 1\}| \\ &= \sum_{\lambda \in \text{Irr}(E)} \epsilon(\lambda) (\lambda(1) + \lambda(u)) - \sum_{\lambda \in \text{Irr}(D)} \epsilon(\lambda) (\lambda(1) + \lambda(u)) \\ &\quad - \sum_{\lambda \in \text{Irr}(E)} \epsilon(\lambda) \lambda(1) + \sum_{\lambda \in \text{Irr}(D)} \epsilon(\lambda) \lambda(1) \\ &= \sum_{\lambda \in \text{Irr}(E)} \epsilon(\lambda) \lambda(u) - \sum_{\lambda \in \text{Irr}(D)} \epsilon(\lambda) \lambda(u) = |\{x \in E \setminus D : x^2 = u\}|. \quad \square \end{aligned}$$

4 Theorems D and E

The following result implies Theorem D.

Theorem 14. *Suppose that B is a real, nilpotent, non-principal 2-block fulfilling the statement of Theorem 13. Then Conjecture B holds for B .*

Proof. Let (D, E) be defect pair of B . By Gow [8, Theorem 5.1], there exists a 2-rational character $\chi_0 \in \text{Irr}(B)$ of height 0 and $\epsilon(\chi_0) = 1$. Let

$$\Gamma : \text{Irr}(D) \rightarrow \text{Irr}(B), \quad \lambda \mapsto \lambda * \chi_0$$

be the Broué–Puig bijection. Let $(u_1, b_1), \dots, (u_k, b_k)$ be representatives for the conjugacy classes of B -subsections. Since B is nilpotent, we may assume that $u_1, \dots, u_k \in D$ represent the conjugacy classes of D . Let $\text{IBr}(b_i) = \{\varphi_i\}$ for $i = 1, \dots, k$. Since χ_0 is 2-rational, we have $\sigma_i := d_{\chi_0, \varphi_i}^u \in \{\pm 1\}$ for $i = 1, \dots, k$. Hence, the generalized decomposition matrix of B has the form

$$Q = (\lambda(u_i)\sigma_i : \lambda \in \text{Irr}(D), i = 1, \dots, k)$$

(see [16, Section 8.10]). Let $v := (\epsilon(\Gamma(\lambda)) : \lambda \in \text{Irr}(D))$ and $w := (w_1, \dots, w_k)$ where $w_i := |\{x \in E \setminus D : x^2 = u_i\}|$. Then Theorem 13 reads as $vQ = w$.

Let $d_i := |\text{C}_D(u_i)|$ and $d = (d_1, \dots, d_k)$. Then the second orthogonality relation yields $Q^t \bar{Q} = \text{diag}(d)$ where Q^t denotes the transpose of Q . It follows that $Q^{-1} = \text{diag}(d)^{-1} \bar{Q}^t$ and

$$v = w \text{diag}(d)^{-1} \bar{Q}^t = w \text{diag}(d)^{-1} Q^t,$$

because $\bar{v} = v$. Since $w_i = |\{x \in E \setminus D : x^2 = u_i^y\}|$ for every $y \in D$, we obtain $\sum_{i=1}^k w_i |D : \text{C}_D(u_i)| = |E \setminus D| = |D|$. In particular,

$$1 = \epsilon(\chi_0) = \sum_{i=1}^k \frac{w_i \sigma_i}{|\text{C}_D(u_i)|} \leq \sum_{i=1}^k \frac{w_i |\sigma_i|}{|\text{C}_D(u_i)|} = 1.$$

Therefore, $\sigma_i = 1$ or $w_i = 0$ for each i . This means that the signs σ_i have no impact on the solution of the linear system $xQ = w$. Hence, we may assume that $Q = (\lambda(u_i))$ is just the character table of D . Since Q has full rank, v is the only solution of $xQ = w$. Setting $\mu(\lambda) := \frac{1}{|D|} \sum_{e \in E \setminus D} \lambda(e^2)$, it suffices to show that $(\mu(\lambda) : \lambda \in \text{Irr}(D))$ is another solution of $xQ = w$. Indeed,

$$\begin{aligned} \sum_{\lambda \in \text{Irr}(D)} \frac{\lambda(u_i)}{|D|} \sum_{e \in E \setminus D} \lambda(e^2) &= \frac{1}{|D|} \sum_{e \in E \setminus D} \sum_{\lambda \in \text{Irr}(D)} \lambda(u_i) \lambda(e^2) \\ &= \frac{1}{|D|} \sum_{\substack{e \in E \setminus D \\ e^2 = u_i^{-1}}} |D : \text{C}_D(u_i)| |\text{C}_D(u_i)| = w_i \end{aligned}$$

for $i = 1, \dots, k$. □

Theorem E. *Conjectures B and C hold for all nilpotent 2-blocks of solvable groups.*

Proof. Let B be a real, nilpotent, non-principal 2-block of a solvable group G with defect pair (D, E) . We first prove Conjecture C for B . Since all sections of G are solvable and all blocks dominated by B -subsections are nilpotent, Conjecture C holds for those blocks as well. Hence, the hypothesis of Theorem 13 is fulfilled for B . Now by Theorem 14, Conjecture B holds for B .

Let $N := \text{O}_{2'}(G)$ and let $\theta \in \text{Irr}(N)$ such that the block $\{\theta\}$ is covered by B . Since B is non-principal, $\theta \neq 1_N$ and therefore $\bar{\theta} \neq \theta$ as N has odd order. Since B also lies over $\bar{\theta}$, it follows that $G_\theta < G$. Let b be the Fong–Reynolds correspondent of B in the extended stabilizer G_θ^* . By [22, Theorem 9.14] and [20, p. 94], the Clifford correspondence $\text{Irr}(b) \rightarrow \text{Irr}(B)$, $\psi \mapsto \psi^G$ preserves decomposition numbers and F-S indicators. Thus, we need to show that b has defect pair (D, E) . Let β be the Fong–Reynolds correspondent of B in G_θ . By [22, Theorem 10.20], β is the unique block over θ . In particular, the block idempotents $1_\beta = 1_\theta$ are the same (we identify θ with the block $\{\theta\}$). Since b is also the unique block of G_θ^* over θ , we have $1_b = 1_\theta + 1_{\bar{\theta}} = \sum_{x \in N} \alpha_x x$ for some $\alpha_x \in F$. Let S be a set of representatives for the cosets G/G_θ^* . Then

$$1_B = \sum_{s \in S} (1_\theta + 1_{\bar{\theta}})^s = \sum_{s \in S} 1_b^s = \sum_{g \in N} \left(\sum_{s \in S} \alpha_{g^{s^{-1}}} \right) g.$$

Hence, there exists a real defect class K of B such that $\alpha_{g^{s^{-1}}} \neq 0$ for some $g \in K$ and $s \in S$. Of course we can assume that $g = g^{s^{-1}}$. Then 1_b does not vanish on g . By [22, Theorem 9.1], the central characters λ_B , λ_b and λ_θ agree on N . It follows that K is also a real defect class of b . Hence, we may assume that (D, E) is a defect pair of b .

It remains to consider $G = G_\theta^*$ and $B = b$. Then D is a Sylow 2-subgroup of G_θ by [22, Theorem 10.20] and E is a Sylow 2-subgroup of G . Since $|G : G_\theta| = 2$, it follows that $G_\theta \trianglelefteq G$ and $N = \text{O}_{2'}(G_\theta)$. By [21, Lemma 1 and 2], β is nilpotent and G_θ is 2-nilpotent, i. e. $G_\theta = N \rtimes D$ and $G = N \rtimes E$. Let $\tilde{\Phi} := \sum_{\chi \in \text{Irr}(B)} \chi(1)\chi = \varphi(1)\Phi$ where $\text{IBr}(B) = \{\varphi\}$. We need to show that

$$\epsilon(\tilde{\Phi}) = \varphi(1) |\{x \in E \setminus D : x^2 = 1\}|.$$

Note that $\chi_N = \frac{\chi(1)}{2\theta(1)}(\theta + \bar{\theta})$. By Frobenius reciprocity, it follows that $\tilde{\Phi} = 2\theta(1)\theta^G$ and

$$\tilde{\Phi}_N = |G : N| \theta(1)(\theta + \bar{\theta}).$$

Since Φ vanishes on elements of even order, $\tilde{\Phi}$ vanishes outside N . Since $\tilde{\Phi}_{G_\theta}$ is a sum of non-real characters in β , we have

$$\epsilon(\tilde{\Phi}) = \frac{1}{|G|} \sum_{g \in G_\theta} \tilde{\Phi}(g^2) + \frac{1}{|G|} \sum_{g \in G \setminus G_\theta} \tilde{\Phi}(g^2) = \frac{1}{|G|} \sum_{g \in G \setminus G_\theta} \tilde{\Phi}(g^2).$$

Every $g \in G \setminus G_\theta = NE \setminus ND$ with $g^2 \in N$ is N -conjugate to a unique element of the form xy where $x \in E \setminus D$ is an involution and $y \in C_N(x)$ (Sylow's theorem). Setting $\Delta := \{x \in E \setminus D : x^2 = 1\}$, we obtain

$$\epsilon(\tilde{\Phi}) = \frac{\theta(1)}{|N|} \sum_{x \in \Delta} |N : C_N(x)| \sum_{y \in C_N(x)} (\theta(y) + \overline{\theta(y)}) = 2\theta(1) \sum_{x \in \Delta} \frac{1}{|C_N(x)|} \sum_{y \in C_N(x)} \theta(y). \quad (5)$$

For $x \in \Delta$ let $H_x := N\langle x \rangle$. Again by Sylow's theorem, the N -orbit of x is the set of involutions in H_x . From $\theta^x = \bar{\theta}$ we see that θ^{H_x} is an irreducible character of 2-defect 0. By [8, Theorem 5.1], we have $\epsilon(\theta^{H_x}) = 1$. Now applying the same argument as before, it follows that

$$1 = \epsilon(\theta^{H_x}) = \frac{1}{|N|} \sum_{g \in H_x \setminus N} \theta^{H_x}(g^2) = \frac{2}{|C_N(x)|} \sum_{y \in C_N(x)} \theta(y).$$

Combined with (5), this yields $\epsilon(\tilde{\Phi}) = 2\theta(1)|\Delta|$. By Green's theorem (see [22, Theorem 8.11]), $\varphi_N = \theta + \bar{\theta}$ and $\epsilon(\tilde{\Phi}) = \varphi(1)|\Delta|$ as desired. \square

For non-principal blocks B of solvable groups with $l(B) = 1$ it is not true in general that G_θ is 2-nilpotent in the situation of Theorem E. For example, a (non-real) 2-block of a triple cover of $A_4 \times A_4$ has a unique simple module. Extending this group by an automorphism of order 2, we obtain the group $G = \text{SmallGroup}(864, 3988)$, which fulfills the assumptions with $D \cong C_2^4$, $N \cong C_3$ and $|G : NE| = 9$.

In order to prove Conjecture C for arbitrary 2-blocks of solvable groups, we may follow the steps in the proof above until E is a Sylow 2-subgroup of G and $|G : G_\theta| = 2$. By [24, Theorem 2.1], one gets

$$\varphi(1)/\theta(1) = 2\sqrt{|G_\theta/N|_{2'}} = \sqrt{|G : EN|}.$$

With some more effort, the claim then boils down to a purely group-theoretical statement:

Let B be a real, non-principal 2-block of a solvable group G with defect pair (D, E) and $l(B) = 1$. Let $N := O_{2'}(G)$ and $\bar{G} := G/N$. Let $\theta \in \text{Irr}(N)$ such $\{\theta\}$ is covered by B . Then

$$|\{\bar{x} \in \bar{G} \setminus \bar{G}_\theta : \bar{x}^2 = 1\}| = |\{x \in E \setminus D : x^2 = 1\}| \sqrt{|G : EN|}.$$

Unfortunately, I am unable to prove this.

Acknowledgment

I thank Gabriel Navarro for providing some arguments for Theorem E from his paper [21]. John Murray and three anonymous referees have made many valuable comments, which improved the quality of the manuscript. The work is supported by the German Research Foundation (SA 2864/3-1 and SA 2864/4-1).

References

- [1] J. An and C. W. Eaton, *Nilpotent Blocks of Quasisimple Groups for the Prime Two*, *Algebr. Represent. Theory* **16** (2013), 1–28.
- [2] R. Brauer, *Representations of finite groups*, in: *Lectures on Modern Mathematics*, Vol. I, 133–175, Wiley, New York, 1963.
- [3] R. Brauer, *Some applications of the theory of blocks of characters of finite groups. III*, *J. Algebra* **3** (1966), 225–255.
- [4] M. Broué and L. Puig, *A Frobenius theorem for blocks*, *Invent. Math.* **56** (1980), 117–128.
- [5] C. W. Eaton, *Generalisations of conjectures of Brauer and Olsson*, *Arch. Math. (Basel)* **81** (2003), 621–626.
- [6] The GAP Group, *GAP – Groups, Algorithms, and Programming, Version 4.12.0*; 2022, (<http://www.gap-system.org>).
- [7] R. Gow, *Real-valued characters and the Schur index*, *J. Algebra* **40** (1976), 258–270.
- [8] R. Gow, *Real-valued and 2-rational group characters*, *J. Algebra* **61** (1979), 388–413.
- [9] R. Gow, *Real 2-blocks of characters of finite groups*, *Osaka J. Math.* **25** (1988), 135–147.

- [10] R. Gow and J. Murray, *Real 2-regular classes and 2-blocks*, J. Algebra **230** (2000), 455–473.
- [11] G. T. Helleloid and U. Martin, *The automorphism group of a finite p -group is almost always a p -group*, J. Algebra **312** (2007), 294–329.
- [12] L. Héthelyi, E. Horváth and E. Szabó, *Real characters in blocks*, Osaka J. Math. **49** (2012), 613–623.
- [13] T. Ichikawa and Y. Tachikawa, *The Super Frobenius–Schur Indicator and Finite Group Gauge Theories on Pin– Surfaces*, to appear in Commun. Math. Phys., DOI: 10.1007/s00220-022-04601-9.
- [14] I. M. Isaacs, *Character theory of finite groups*, AMS Chelsea Publishing, Providence, RI, 2006.
- [15] R. Kessar and G. Malle, *Quasi-isolated blocks and Brauer’s height zero conjecture*, Ann. of Math. (2) **178** (2013), 321–384.
- [16] M. Linckelmann, *The block theory of finite group algebras. Vol. II*, London Mathematical Society Student Texts, Vol. 92, Cambridge University Press, Cambridge, 2018.
- [17] J. Murray, *Strongly real 2-blocks and the Frobenius–Schur indicator*, Osaka J. Math. **43** (2006), 201–213.
- [18] J. Murray, *Components of the involution module in blocks with cyclic or Klein-four defect group*, J. Group Theory **11** (2008), 43–62.
- [19] J. Murray, *Real subpairs and Frobenius–Schur indicators of characters in 2-blocks*, J. Algebra **322** (2009), 489–513.
- [20] J. Murray, *Frobenius–Schur indicators of characters in blocks with cyclic defect*, J. Algebra **533** (2019), 90–105.
- [21] G. Navarro, *Nilpotent characters*, Pacific J. Math. **169** (1995), 343–351.
- [22] G. Navarro, *Characters and blocks of finite groups*, London Mathematical Society Lecture Note Series, Vol. 250, Cambridge University Press, Cambridge, 1998.
- [23] G. Navarro, *Character theory and the McKay conjecture*, Cambridge Studies in Advanced Mathematics, Vol. 175, Cambridge University Press, Cambridge, 2018.
- [24] G. Navarro, B. Späth and P. H. Tiep, *On fully ramified Brauer characters*, Adv. Math. **257** (2014), 248–265.
- [25] S. Trefethen and C. R. Vinroot, *A computational approach to the Frobenius–Schur indicators of finite exceptional groups*, Internat. J. Algebra Comput. **30** (2020), 141–166.
- [26] W. Willems, *Duality and forms in representation theory*, in: Representation theory of finite groups and finite-dimensional algebras (Bielefeld, 1991), 509–520, Progr. Math., Vol. 95, Birkhäuser, Basel, 1991.